arXiv:1303.3144v1 [gr-qc] 13 Mar 2013

On the Gravitational Energy-Momentum Vector in f(T) Theories

S. C. Ulhoa[*]

Faculdade Gama,Universidade de Brasília, 72444-240,
Setor Leste (Gama), Brasília, DF, Brazil.

E. P. Spaniol[†]

UDF Centro Universitário and Centro Universitário IESB
(Campus Sul Edson Machado), Brasília, DF, Brazil.

(Dated: March 14, 2013)

Abstract

This work is devoted to analyze the energy of the Universe in the context of f(T) theories. Such theories are the analogous counterpart of the well known f(R) theories that, however, uses torsion instead of curvature. We obtain a general expression for the gravitational energy-momentum vector in this framework. Using the hypothesis of the isotropy of spacetime, we find the energy of the Universe and compare it with the energy obtained in the realm of teleparallelism equivalent to general relativity (TEGR).

PACS numbers: 04.20-q; 04.20.Cv; 02.20.Sv

Keywords: Modified Gravity; Energy-momentum.

---

[*] sc.ulhoa@gmail.com

[†] spaniol.ep@gmail.com

I. INTRODUCTION

Teleparallel gravity and general relativity are two different theories with the same field equations. In general relativity it is not possible to construct an expression for the gravitational energy in terms of the metric tensor and its second derivatives which gave rise the the approach of pseudo-tensors of energy-momentum. On the other hand in TEGR there is an expression for the gravitational energy-momentum vector, tested along the years, in terms of the tetrad field which is the dynamical variable of the theory. This feature will be explored in this article.

The Hilbert-Einstein lagrangian density, which gives the dynamics of general relativity, can be generalized as a function of Ricci scalar. This establishes a wide class of lagrangian densities that lead to what is known as f(R) theories [1]. Several cosmological observations, such as supernovae [2, 3], cosmic microwave background [4, 5], large-scale structure [6, 7] and baryon acoustic oscillations [8], point to exotic kind of energy which is known as dark energy. The f(R) theories have been used successfully to explain such observations. It can explain the experimental data since it adds extra terms in the energy-momentum tensor of matter fields in the Einstein equations, working as a different font of energy density.

Such a success leads us to study the generalization of the lagrangian density of teleparallel gravity which is known as f(T) theories. Recently, models based on f(T) gravity were presented as an alternative to inflationary models [9]. Moreover, in the literature we find some works that attempt to explain the late-time accelerated expansion of the Universe in the context of f(T) theories [10, 11]. We would like to shed light on the subject by exploring the main advantage of f(T) theories: the existence of a gravitational energy-momentum vector. In our opinion dark energy could be a manifestation of some sort of gravitational energy, hence studying it in the framework of f(T) theories become rather an imposition. Therefore the knowledge about the energy of the Universe can improve our understanding of how it is expanding.

Recently the energy of the Universe was obtained in the context of teleparallel gravity [12]. Such an result was derived considering the Universe as described by the FRW metric and the use of regularized expressions as well. This use is necessary in cosmology to describe reference frames since it is possible to have remanence of torsion in the flat space-time. Thus we intend to find a general expression of the energy-momentum vector in the context of f(T) theories, contained in an arbitrary volume V of the three-dimensional spacelike hypersurface, P^a. The definition of P^a proved to be invariant under coordinate transformation and transform like a vector under Lorentz transformations, features that are essential to a true energy-momentum vector. Then, as an application we obtain the gravitational energy of the universe taking into account the cosmological principle and analyze what would be a correction in the gravitational energy from the approach of the f(T) theories when compared to TEGR.

Notation: space-time indices $\mu, \nu, ...$ and SO(3,1) indices $a, b, ...$ run from 0 to 3. Time and space indices are indicated according to $\mu = 0, i, \quad a = (0), (i)$. The tetrad field is denoted by $e^a{}_\mu$ and the determinant of the tetrad field is represented by $e = \det(e^a{}_\mu)$.

II. TELEPARALLEL GRAVITY AND F(T) THEORIES

In this section we would like to establish the field equations of f(T) theories and a definition of energy-momentum vector. However firstly it is necessary to recall some ideas of TEGR which is an alternative theory of gravitation, that is entirely equivalent to standard general relativity in what concerns dynamical equations. The main difference between both formulations is the existence, in the framework of Teleparallel gravity, of a true gravitational energy-momentum vector which is independent of coordinate transformations and sensible to the change of reference frame. It replaces the Riemannian curvature by the torsion in a tetrad formulation of Weitzenböck (or Cartan) space-time [13, 14]. An approach originally considered by Einstein himself in 1930 [15].

The tetrad field and metric tensor are related by $g^{\mu\nu} = e^{a\mu} e_a{}^\nu$. The familiar theory of general relativity deals with the Christoffel symbols ${}^0\Gamma_{\mu\lambda\nu}$, as the connection of space-time. On the other hand TEGR is formulated in terms of Cartan connection [16], $\Gamma_{\mu\lambda\nu} = e^a{}_\mu \partial_\lambda e_{a\nu}$. The geometric framework of both theories is related by means the following identity

$$\Gamma_{\mu\lambda\nu} = {}^0\Gamma_{\mu\lambda\nu} + K_{\mu\lambda\nu} \,, \tag{1}$$

where $K_{\mu\lambda\nu}$ is given by

$$K_{\mu\lambda\nu} = \frac{1}{2}(T_{\lambda\mu\nu} + T_{\nu\lambda\mu} + T_{\mu\lambda\nu}) \,. \tag{2}$$

$K_{\mu\lambda\nu}$ is the contortion tensor defined in terms of the torsion tensor constructed from the Cartan connection. The torsion tensor is $T_{\mu\lambda\nu} = e_{a\mu}T^a{}_{\lambda\nu}$, with

$$T^a{}_{\lambda\nu} = \partial_\lambda e^a{}_\nu - \partial_\nu e^a{}_\lambda \,. \tag{3}$$

The curvature tensor obtained from $\Gamma_{\mu\lambda\nu}$ is identically zero. From the identity (1) we have

$$eR(e) \equiv -e(\frac{1}{4}T^{abc}T_{abc} + \frac{1}{2}T^{abc}T_{bac} - T^aT_a) + 2\partial_\mu(eT^\mu)\,, \tag{4}$$

where $R(e)$ is the scalar curvature of a Riemannian manifold in terms of the tetrad field and $T^\mu = T^b{}_b{}^\mu$.

The Teleparallel Lagrangian density can be defined from the identity (4) which reads

$$\begin{aligned} \mathfrak{L}(e_{a\mu}) &= -\kappa\, e\,(\frac{1}{4}T^{abc}T_{abc} + \frac{1}{2}T^{abc}T_{bac} - T^aT_a) - \mathfrak{L}_M \\ &\equiv -\kappa\, e\Sigma^{abc}T_{abc} - \mathfrak{L}_M\ , \end{aligned} \tag{5}$$

where $\kappa = 1/(16\pi)$, $\mathfrak{L}_M$ is the Lagrangian density of matter fields and Σ^{abc} is given by

$$\Sigma^{abc} = \frac{1}{4}(T^{abc} + T^{bac} - T^{cab}) + \frac{1}{2}(\eta^{ac}T^b - \eta^{ab}T^c)\ , \tag{6}$$

with $T^a = e^a{}_\mu T^\mu$. It is important to note that the divergence term has been dropped once it does not contribute to the field equations. Thus both theories share the same field equations and hence are equivalent to each other, since there is an equivalence between their Lagrangian densities.

The most general Lagrangian density in the realm of teleparallelism is given by

$$\mathfrak{L} = e\, f(T) - \mathfrak{L}_M\,, \tag{7}$$

where $T = \Sigma^{abc}T_{abc}$. Performing a variational derivative of the above Lagrangian density with respect to $e_{a\gamma}$, which are the dynamical variables of the system, the field equations are

$$f'(T)\left[\partial_\nu\left(e\,\Sigma^{a\gamma\nu}\right) - e\,\Sigma^{bc\gamma}T_{bc}{}^a\right] - e\,\Sigma^{a\lambda\gamma}\left(\partial_\lambda T\right) f''(T) + \frac{1}{4}e\, e^{a\gamma} f(T) = \frac{1}{4\kappa}\, e\, T^{a\gamma}\,, \tag{8}$$

where $T^{a\mu} = e^a{}_\lambda T^{\mu\lambda} = \frac{1}{e}\frac{\delta\mathcal{L}_M}{\delta e_{a\mu}}$ is the energy-momentum tensor of matter fields. The prime in $f(T)$ means a derivative with respect to T. The field equations can be rewritten as

$$\partial_\nu\left(e\Sigma^{a\lambda\nu}\, f'(T)\right) = \frac{1}{4\kappa} e\, e^a{}_\mu (t^{\lambda\mu} + T^{\lambda\mu})\,, \tag{9}$$

where $t^{\lambda\mu}$ is defined by

$$t^{\lambda\mu} = \kappa\left[4\, f'(T)\, \Sigma^{bc\lambda} T_{bc}{}^{\mu} - g^{\lambda\mu}\, f(T)\right]\,. \tag{10}$$

Since $\Sigma^{a\lambda\nu}$ is skew-symmetric in the last two indices, it follows that

$$\partial_\lambda\partial_\nu\left(e\Sigma^{a\lambda\nu}\, f'(T)\right) \equiv 0\,. \tag{11}$$

Thus we get

$$\partial_\lambda(et^{a\lambda} + eT^{a\lambda}) = 0 \tag{12}$$

which yields the continuity equation

$$\frac{d}{dt}\int_V d^3x\, e\, e^a{}_\mu (t^{0\mu} + T^{0\mu}) = -\oint_S dS_j\, \left[e\, e^a{}_\mu (t^{j\mu} + T^{j\mu})\right]\,.$$

It should be noted that the above expression works as a conservation law for the sum of energy-momentum tensor of matter fields and for the quantity $t^{\lambda\mu}$. Thus $t^{\lambda\mu}$ is interpreted as the energy-momentum tensor of the gravitational field in the context of $f(T)$ theories, being more general than (and slightly different from) the usual quantity of TEGR [17], [18]. Therefore, one can write the total energy-momentum contained in a three-dimensional volume V of space as

$$P^a = \int_V d^3x\, e\, e^a{}_\mu (t^{0\mu} + T^{0\mu})\,, \tag{13}$$

or using the field equations we have

$$P^a = 4k\int_V d^3x\, \partial_\nu\left(e\,\Sigma^{a0\nu}\, f'(T)\right)\,. \tag{14}$$

It is worth to note that the above expression is invariant under coordinate transformation and transforms like a vector under Lorentz transformations. Such features are desirable and expected for a true energy-momentum vector.

III. APPLICATION IN COSMOLOGICAL SCALES: THE ENERGY OF THE UNIVERSE

The cosmological principle asserts that the large-scale structure of the Universe reveals homogeneity and isotropy [19]. The most general form of a line element that preserves such features may be written as [20],

$$ds^2 = -dt^2 + a^2(t)\left[\frac{dr^2}{(1-k'r^2)} + r^2(d\theta^2 + \sin^2\theta d\phi^2)\right], \tag{15}$$

where $a(t) = S(t)/|K|^{1/2}$ if $K \neq 0$ and $a(t) = S(t)$ if $K = 0$. $S(t)$ is the scale factor and K is the constant curvature of space. Here $K = |K|k'$ where k' assumes the values $+1, 0, -1$ which correspond to a space of constant positive curvature, a flat space or a space of constant negative curvature, respectively.

Let us choose the following tetrad field adapted to a stationary reference frame

$$e^a{}_\mu = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \sqrt{g_{11}}\sin\theta\cos\phi & \sqrt{A}\cos\theta\cos\phi & -\sqrt{A}\sin\theta\sin\phi \\ 0 & \sqrt{g_{11}}\sin\theta\sin\phi & \sqrt{A}\cos\theta\sin\phi & \sqrt{A}\sin\theta\cos\phi \\ 0 & \sqrt{g_{11}}\cos\theta & -\sqrt{A}\sin\theta & 0 \end{bmatrix} \tag{16}$$

where $A = r^2a^2$. By direct inspection one can see that the above tetrad yields the line element (15). The non-vanishing components of the torsion tensor are

$$\begin{aligned} T_{101} &= \frac{1}{2}\left(\frac{\partial g_{11}}{\partial t}\right), \\ T_{202} &= \frac{1}{2}\left(\frac{\partial A}{\partial t}\right), \\ T_{212} &= \frac{1}{2}\left[\left(\frac{\partial A}{\partial r}\right) - 2\sqrt{g_{11}\,A}\right], \\ T_{303} &= \frac{1}{2}\left(\frac{\partial A}{\partial t}\right)\sin^2\theta\,, \\ T_{313} &= \frac{1}{2}\left[\left(\frac{\partial A}{\partial r}\right) - 2\sqrt{g_{11}\,A}\right]\sin^2\theta\,. \end{aligned} \tag{17}$$

After some algebraic manipulation we find

$$e\,\Sigma^{(0)01} = -\frac{1}{2}\left[\frac{\left(\frac{\partial A}{\partial r}\right) - 2\sqrt{g_{11}}\,A}{\sqrt{g_{11}}}\right]\sin\theta \tag{18}$$

which is simplified to

$$e\,\Sigma^{(0)01} = \left[1 - \left(1 - k'\,r^2\right)^{1/2}\right] ra\sin\theta\,. \tag{19}$$

Our intent when writing such a component is to give the gravitational energy which is the zero component of the vector defined in (14). Therefore we also have

$$T = \frac{1}{2}\left[\frac{-\left(\frac{\partial A}{\partial r}\right)^2 + 4\sqrt{g_{11}A}\left(\frac{\partial A}{\partial r}\right) + 2\left(\frac{\partial A}{\partial t}\right)\left(\frac{\partial g_{11}}{\partial t}\right)A + \left(\frac{\partial A}{\partial t}\right)^2 g_{11} - 4\,g_{11}A}{A^2 g_{11}}\right], \tag{20}$$

which reads

$$T = 6H^2 - \frac{4}{r^2a^2} + 2\,\frac{k'}{a^2} + 4\,\frac{\left(1 - k'\,r^2\right)^{1/2}}{r^2a^2} \tag{21}$$

with $H = \dot{a}/a$ being the Hubble's constant. Let us choose the following form for $f(T)$

$$f(T) = T + \frac{\lambda}{2}T^2$$

where λ is a coupling constant; its value should be very small when compared to T, since there is no observational data concerning modifications of general relativity. Thus $f'(T) = 1 + \lambda T$ and the energy is written as

$$\begin{aligned} P^{(0)} = 4k \lim_{r\to R}\int_0^{2\pi}\int_0^{\pi}&\left[1 - \left(1 - k'\,r^2\right)^{1/2}\right]\cdot \\ &\cdot\left[1 + 2\lambda\left(3H^2 - \frac{2}{r^2a^2} + \frac{k'}{a^2} + \frac{2\left(1 - k'\,r^2\right)^{1/2}}{r^2a^2}\right)\right] ra\sin\theta d\theta d\phi\,, \end{aligned}$$

which becomes

$$P^{(0)} = Ra\left[1 - \left(1 - k'\,R^2\right)^{1/2}\right]\left\{1 + \lambda\left[6H^2 + 2\,\frac{k'}{a^2} - \frac{4}{R^2a^2} + 4\,\frac{\left(1 - k'\,R^2\right)^{1/2}}{R^2a^2}\right]\right\}, \tag{22}$$

after integration over a spherical hyper-surface of radius R.

The quantity $\tilde{P}^{(0)} = Ra\left[1 - (1 - k'\,R^2)^{1/2}\right]$ is precisely the energy of the Universe obtained in the context of TEGR by means of a regularization procedure (see eq. (31) of [12]).

Therefore the correction in the energy of the Universe $\Delta P^{(0)} = P^{(0)} - \tilde{P}^{(0)}$, if the dynamics was modified by the lagrangian density (7), is

$$\Delta P^{(0)} = \lambda \tilde{P}^{(0)} \left[6H^2 + 2\,\frac{k'}{a^2} - \frac{4}{R^2 a^2} + 4\,\frac{\left(1 - k'\,R^2\right)^{1/2}}{R^2 a^2} \right] . \tag{23}$$

It should be noted that both the correction $\Delta P^{(0)}$ and the energy without modification $\tilde{P}^{(0)}$ vanish when the curvature of the Universe is zero since they are proportional to each other. For the two other possible values of k', the extra amount of energy given by eq. (23) could play an important role in what is called dark energy.

The temporal evolution of $a(t)$ is given by the equations

$$3\left(H^2 + \frac{k'}{a^2}\right) + \lambda\left(27H^2 + 30\frac{k'H^2}{a^2} + 7\frac{k'^2}{a^4}\right) = 8\pi\rho\,, \tag{24}$$

$$\frac{\ddot{a}}{a} + 2H^2 + \frac{k'}{a^2} + \lambda\left[18H^2\frac{\ddot{a}}{a} + \frac{6k'}{a^2}H^2 + \frac{2k'}{a^3}\left(\ddot{a} + \frac{k'}{a}\right)\right] = -4\pi p\,, \tag{25}$$

where ρ and p are the energy density and the pressure of the cosmological fluid respectively. Let us reorganize both equations above as

$$3\left(H^2 + \frac{k'}{a^2}\right) = 8\pi\rho_T\,, \tag{26}$$

$$\frac{\ddot{a}}{a} + 2H^2 + \frac{k'}{a^2} + = -4\pi p_T\,, \tag{27}$$

where

$$\rho_T = \rho - \frac{\lambda}{8\pi}\left(27H^2 + 30\frac{k'H^2}{a^2} + 7\frac{k'^2}{a^4}\right)\,,$$

and

$$p_T = p + \frac{\lambda}{4\pi}\left[\frac{18\ddot{a}}{a}H^2 + \frac{6k'}{a^2}H^2 + \frac{2k'}{a^3}\left(\ddot{a} + \frac{k'}{a}\right)\right]\,.$$

It should be noted that the equation of state has been obtained considering the isotropy and homogeneity principle which asserts that any direction is privileged in the Universe and there is a uniform distribution of energy/matter on large scales. Then we drop out any term which decays like r^{-1} or more.

IV. CONCLUSION

In this paper we have obtained a general expression for the gravitational energy-momentum in the realm of teleparallel gravity when the dynamics is governed by a general lagrangian as a function of the scalar $T = \Sigma^{abc}T_{abc}$ and thus called $f(T)$ theories (in analogy to the $f(R)$ theories). Such an expression has never appeared in the literature. Then we applied our expression to the FRW metric which was established taking into account the principle of isotropy and homogeneity of the large scale of the Universe. We suppose a form for the function $f(T)$ and we obtained a correction, $\Delta P^{(0)}$, in the gravitational energy of the Universe. Such a correction can be viewed as a possible explanation for what is known as dark energy since the equation of state for $a(t)$ is modified due to the alteration the lagrangian. Therefore this extra amount of gravitational energy could be seen as a source for dark energy.

[1] Antonio De Felice and Shinji Tsujikawa. f(r) theories. *Living Reviews in Relativity*, 13(3), 2010.

[2] S. Perlmutter, G. Aldering, G. Goldhaber, R. A. Knop, P. Nugent, P. G. Castro, S. Deustua, S. Fabbro, A. Goobar, D. E. Groom, I. M. Hook, A. G. Kim, M. Y. Kim, J. C. Lee, N. J. Nunes, R. Pain, C. R. Pennypacker, R. Quimby, C. Lidman, R. S. Ellis, M. Irwin, R. G. McMahon, P. Ruiz-Lapuente, N. Walton, B. Schaefer, B. J. Boyle, A. V. Filippenko, T. Matheson, A. S. Fruchter, N. Panagia, H. J. M. Newberg, W. J. Couch, and The Supernova Cosmology Project. Measurements of and from 42 high-redshift supernovae. *The Astrophysical Journal*, 517(2):565, 1999.

[3] Adam G. Riess, Alexei V. Filippenko, Peter Challis, Alejandro Clocchiatti, Alan Diercks, Peter M. Garnavich, Ron L. Gilliland, Craig J. Hogan, Saurabh Jha, Robert P. Kirshner, B. Leibundgut, M. M. Phillips, David Reiss, Brian P. Schmidt, Robert A. Schommer, R. Chris Smith, J. Spyromilio, Christopher Stubbs, Nicholas B. Suntzeff, and John Tonry. Observational evidence from supernovae for an accelerating universe and a cosmological constant. *The Astronomical Journal*, 116(3):1009, 1998.

[4] D. N. Spergel, L. Verde, H. V. Peiris, E. Komatsu, M. R. Nolta, C. L. Bennett, M. Halpern,

G. Hinshaw, N. Jarosik, A. Kogut, M. Limon, S. S. Meyer, L. Page, G. S. Tucker, J. L. Weiland, E. Wollack, and E. L. Wright. First-year wilkinson microwave anisotropy probe (wmap) observations: Determination of cosmological parameters. *The Astrophysical Journal Supplement Series*, 148(1):175, 2003.

[5] D. N. Spergel, R. Bean, O. Dor, M. R. Nolta, C. L. Bennett, J. Dunkley, G. Hinshaw, N. Jarosik, E. Komatsu, L. Page, H. V. Peiris, L. Verde, M. Halpern, R. S. Hill, A. Kogut, M. Limon, S. S. Meyer, N. Odegard, G. S. Tucker, J. L. Weiland, E. Wollack, and E. L. Wright. Three-year wilkinson microwave anisotropy probe (wmap) observations: Implications for cosmology. *The Astrophysical Journal Supplement Series*, 170(2):377, 2007.

[6] Max Tegmark, Michael A. Strauss, Michael R. Blanton, Kevork Abazajian, Scott Dodelson, Havard Sandvik, Xiaomin Wang, David H. Weinberg, Idit Zehavi, Neta A. Bahcall, Fiona Hoyle, David Schlegel, Roman Scoccimarro, Michael S. Vogeley, Andreas Berlind, Tamás Budavari, Andrew Connolly, Daniel J. Eisenstein, Douglas Finkbeiner, Joshua A. Frieman, James E. Gunn, Lam Hui, Bhuvnesh Jain, David Johnston, Stephen Kent, Huan Lin, Reiko Nakajima, Robert C. Nichol, Jeremiah P. Ostriker, Adrian Pope, Ryan Scranton, Uro š Seljak, Ravi K. Sheth, Albert Stebbins, Alexander S. Szalay, István Szapudi, Yongzhong Xu, James Annis, J. Brinkmann, Scott Burles, Francisco J. Castander, Istvan Csabai, Jon Loveday, Mamoru Doi, Masataka Fukugita, Bruce Gillespie, Greg Hennessy, David W. Hogg, Željko Ivezić, Gillian R. Knapp, Don Q. Lamb, Brian C. Lee, Robert H. Lupton, Timothy A. McKay, Peter Kunszt, Jeffrey A. Munn, Liam O'Connell, John Peoples, Jeffrey R. Pier, Michael Richmond, Constance Rockosi, Donald P. Schneider, Christopher Stoughton, Douglas L. Tucker, Daniel E. Vanden Berk, Brian Yanny, and Donald G. York. Cosmological parameters from sdss and wmap. *Phys. Rev. D*, 69:103501, May 2004.

[7] Uro š Seljak, Alexey Makarov, Patrick McDonald, Scott F. Anderson, Neta A. Bahcall, J. Brinkmann, Scott Burles, Renyue Cen, Mamoru Doi, James E. Gunn, Željko Ivezić, Stephen Kent, Jon Loveday, Robert H. Lupton, Jeffrey A. Munn, Robert C. Nichol, Jeremiah P. Ostriker, David J. Schlegel, Donald P. Schneider, Max Tegmark, Daniel E. Vanden Berk, David H. Weinberg, and Donald G. York. Cosmological parameter analysis including sdss α forest and galaxy bias: Constraints on the primordial spectrum of fluctuations, neutrino mass, and dark energy. *Phys. Rev. D*, 71:103515, May 2005.

[8] Daniel J. Eisenstein, Idit Zehavi, David W. Hogg, Roman Scoccimarro, Michael R. Blan-

ton, Robert C. Nichol, Ryan Scranton, Hee-Jong Seo, Max Tegmark, Zheng Zheng, Scott F. Anderson, Jim Annis, Neta Bahcall, Jon Brinkmann, Scott Burles, Francisco J. Castander, Andrew Connolly, Istvan Csabai, Mamoru Doi, Masataka Fukugita, Joshua A. Frieman, Karl Glazebrook, James E. Gunn, John S. Hendry, Gregory Hennessy, Zeljko Ivezi, Stephen Kent, Gillian R. Knapp, Huan Lin, Yeong-Shang Loh, Robert H. Lupton, Bruce Margon, Timothy A. McKay, Avery Meiksin, Jeffery A. Munn, Adrian Pope, Michael W. Richmond, David Schlegel, Donald P. Schneider, Kazuhiro Shimasaku, Christopher Stoughton, Michael A. Strauss, Mark SubbaRao, Alexander S. Szalay, Istvn Szapudi, Douglas L. Tucker, Brian Yanny, and Donald G. York. Detection of the baryon acoustic peak in the large-scale correlation function of sdss luminous red galaxies. *The Astrophysical Journal*, 633(2):560, 2005.

[9] Rafael Ferraro and Franco Fiorini. Modified teleparallel gravity: Inflation without an inflaton. *Phys. Rev. D*, 75:084031, Apr 2007.

[10] Gabriel R. Bengochea and Rafael Ferraro. Dark torsion as the cosmic speed-up. *Phys. Rev. D*, 79:124019, Jun 2009.

[11] Eric V. Linder. Einstein's other gravity and the acceleration of the universe. *Phys. Rev. D*, 81:127301, Jun 2010.

[12] S.C. Ulhoa, J.F da Rocha Neto, and J.W. Maluf. The Gravitational Energy Problem for Cosmological Models in Teleparallel Gravity. *Int.J.Mod.Phys.*, D19:1925–1935, 2010.

[13] Kenji Hayashi and Takeshi Shirafuji. New general relativity. *Phys. Rev. D*, 19:3524–3553, Jun 1979.

[14] José W. Maluf. Hamiltonian formulation of the teleparallel description of general relativity. *Journal of Mathematical Physics*, 35(1):335–343, 1994.

[15] A. Einstein. Unified field theory based on riemannian metrics and distant parallelism. *Math. Annal.*, 102:685–697, 1930.

[16] E. Cartan. On a Generalization of the Notion of Reimann Curvature and Spaces with Torsion. In P. G. Bergmann and V. de Sabbata, editors, *NATO ASIB Proc. 58: Cosmology and Gravitation: Spin, Torsion, Rotation, and Supergravity*, pages 489–491, 1980.

[17] J. W. Maluf. The gravitational energy-momentum tensor and the gravitational pressure. *Annalen Phys.*, 14:723–732, 2005.

[18] V. C. de Andrade, L. C. T. Guillen, and J. G. Pereira. Gravitational energy-momentum density in teleparallel gravity. *Phys. Rev. Lett.*, 84:4533–4536, May 2000.

[19] S. Weinberg. *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity.* John and Wiley & Sons, Inc., 1972.

[20] Ray d'Inverno. *Introducing Einstein's Relativity.* Clarendon Press, Oxford, 4th edition, 1996.